

002250-197

UNITED STATES PLANT PATENT APPLICATION

of

PETER P. LIM

and

MARTIN NEMKO

for

SHRUB ROSE PLANT NAMED 'BAIPOME'

BOTANICAL COMMERCIAL CLASSIFICATION

Rosa hybrida/Shrub Rose Plant

VARIETAL DENOMINATION

cv. 'BAIpome'

Summary of the Invention

The new variety of *Rosa hybrida* landscape shrub rose plant was created at Yamhill, Oregon during 1995 by artificial pollination wherein two parents were crossed which previously had been studied in the hope that they would contribute the desired characteristics. The female parent (*i.e.*, the seed parent) was the 'Noatraum' variety (United States Plant Patent No. 7,282). The male parent (*i.e.*, the pollen parent) also was the 'Noatraum' variety. The 'Noatraum' variety was bred by Werner Noack of Guetersloh, Germany. The parentage of the new variety can be summarized as follows:

'Noatraum' × 'Noatraum'.

The seeds resulting from the above pollination were sown and plants were obtained which were physically and biologically different from each other. Selective study resulted in the identification during 1998 of a single plant of the new variety having the distinctive ornamental characteristics hereafter described.

It was found that the new variety of the present invention:

- (a) exhibits a compact, tight, and rounded growth habit,
- (b) abundantly forms on a continuous basis in clusters attractive single cup-shaped light soft pink blossoms having a white center,
- (c) forms glossy medium green adult foliage,
- (d) displays excellent tolerance to Black Spot and Powdery Mildew; and
- (e) is particularly well suited for growing as attractive ornamentation in parks and gardens.

The young rose pink flower buds open to the light pink blossoms having a white center that cover the bush throughout the season. The aging petals commonly detach cleanly and do not detract from the attractiveness of the blooms. A tough shrub rose plant having good hardiness is provided.

The new variety is a small stature versatile landscape plant that well meets the needs of the horticultural industry. It can be grown to advantage anywhere in the landscape where attractive ornamentation is desired. The distinctive blossoms are displayed throughout the season. The plant can be grown as a specimen, border planting, or as a mass ground cover.

The new variety can be readily distinguished from other varieties including its parental 'Noatraum' variety. More specifically, the 'Noatraum' variety forms much darker pink blossoms with near white centers.

The new variety has been found to undergo asexual propagation by budding on an understock and by the rooting of cuttings. 'Dr. Huey' rootstock was used during the budding. Such asexual propagation by these techniques at Yamhill, Oregon, St. Paul, Minnesota, and at Phoenix, Arizona has shown that the characteristics of the new variety are stable and are strictly transmissible by such asexual propagation from one generation to another. The new variety grows well on its own roots.

Initially the new plant was designated 95K61-1P. The new variety subsequently has been named the 'BAIpome' variety, and currently is being marketed while using the PINK GNOME and GARDEN PATH trademarks.

Brief Description of the Photographs

The accompanying photographs show as nearly true as it is reasonably possible to make the same in a color illustrations of this character, typical specimens of the new variety. The rose plants of the new variety had been propagated by the rooting of cuttings, were approximately two and three years of age, and were being grown outdoors at St. Paul, Minnesota and at Yamhill, Oregon.

FIG. 1 illustrates typical plants with blossoms in various stages of opening and foliage of the new variety. Clusters of the attractive single cup-shaped soft pink blossoms with white centers are shown.

FIG. 2 illustrates a close view of typical buds and fully open flowers of the new variety in greater detail.

Detailed Description

The chart used in the identification of the colors is that of the Royal Horticultural Society (R.H.S. Colour Chart). Common color terms are to be accorded their ordinary dictionary significance. The description is based on the observation of two and three year-old plants while grown outdoors on their own roots during June and July 2002 and 2003 at St. Paul, Minnesota, and July 2003 at Yamhill, Oregon.

BOTANICAL CLASSIFICATION: *Rosa hybrida*, var. 'BAIpome'.

CLASS: Landscape Shrub.

PLANT:

<i>height</i>	--	approximately 30 cm on average at the end of the growing season.
<i>width</i>	--	approximately 50 cm on average at the end of the growing season.
<i>habit</i>	—	compact, tight, and rounded.

BRANCHES:

- color* -- *young stems*: Green Group 138B with highlights of Greyed-Purple Group 183C.
- *adult wood*: commonly between Green Group 137B and 137C.
- thorns* -- *size*: commonly approximately 0.4 cm in length on average with prickles on the younger stems.
- *color*: commonly between Greyed-Purple Group 187B and 187C.

LEAVES:

- stipules* -- generally parallel with the auricle facing outward.
- petioles* -- *upper surface*: Green Group 137B with a center valley of Green Group 138C.
- *under surface*: Green Group 138B.
- leaflets* -- *number*: 3, 5 (most often), and 7.
- *shape*: ovate with a rounded base and an acuminate tip.
- *serration*: serrulate.

- *texture*: bears a glossy finish and is physically firm.
- *general appearance*: medium green with a glossy surface when mature.
- *color (young foliage)*:
upper surface: Green Group 137B.
under surface: Green Group 137C.
- *color (adult foliage)*:
upper surface: Green Group 137B and 137C.
under surface: commonly between Green Group 137D and 138B.

INFLORESCENCE

- number of flowers* -- commonly approximately 18 blooms on average per cluster on a lateral stem.
- peduncle* -- Yellow-Green Group 144A with highlights of Red-Purple Group 59A in coloration, and the length is approximately 1.85 cm on average.

- sepals*
- *upper surface*: Yellow-Green Group 146B
fading at the margin to Yellow-Green
Group 144C.
 - *under surface*: Yellow-Green Group 146C
fading at the margin to Yellow-Green
Group 145C.
 - *size*: approximately 0.6 cm in length.
- buds*
- *shape*: generally rounded.
 - *size*: small.
 - *length*: approximately 0.75 cm on average.
 - *diameter*: approximately 0.75 cm on
average.
 - *color*:
upper surface: commonly between
Red-Purple Group 58C and 58D fading to
Yellow Group 2B at the base.
under surface: Red-Purple Group 58C with
some Red-Purple Group 62A particularly at
the veins.
- flower*
- *shape*: single and somewhat informal.
 - *diameter*: approximately 2 cm on average.

-- *color (when opening begins):*

upper surface: between Red-Purple Group 62A and 62B with highlights of Red-Purple Group 63C at the margin.

under surface: Red-Purple Group 62A with highlights of Red-Purple Group 63C.

-- *color (when blooming):*

upper surface: Red-Purple Group 62B blending to White Group 155C at the center.

under surface: Red-Purple Group 62C blending to White Group 155D at the center.

-- *color (at end of blooming):*

upper surface: Red-Purple Group 62D over approximately 65 percent of the surface and blending to White Group 155D at the center.

under surface: Red-Purple Group 62D over approximately 65 percent of the surface and blending to White Group 155D at the center.

- *color (basal petal spot):*
White Group 155A.
- *fragrance:* slight green apple.
- *lasting quality:* the blossoms commonly
last approximately 5 to 7 days on average
depending upon the temperature and
sunlight exposure.
- *petal number:* 5 under normal growing
conditions.
- *petal shape:* heart-shaped.
- *petal drop:* commonly the petals detach
cleanly at full blossom maturity.
- *anthers:* Greyed-Orange Group 163B in
coloration.
- *filament number:* commonly approximately
58 on average.
- *filament color:* Greyed-Yellow Group
162A in coloration.
- *pistils:* approximately 20 on average.
- *stigmas:* Yellow-Green Group 150A in
coloration.
- *styles:* Yellow Group 7A in coloration.

- *receptacle*: separate and free.
- *hip shape*: oval.
- *hip length*: approximately 1.0 cm.
- *hip width*: approximately 7.0 mm.
- *seeds*: medium beige in coloration, and approximately 4.0 mm in length.

DEVELOPMENT

- | | | |
|-------------------------------|----|--------------------------------------------------------|
| <i>vegetation</i> | -- | vigorous, and compact. |
| <i>blooming</i> | -- | abundant and continuous throughout the season. |
| <i>resistance to diseases</i> | -- | excellent tolerance to Black Spot, and Powdery Mildew. |
| <i>hardiness zone</i> | -- | U.S.D.A. Hardiness Zone Nos. 4 to 9. |